

# ON THE FEICHTINGER CONJECTURE

P. GĂVRUȚA

**ABSTRACT.** We prove the Feichtinger Conjecture for a class of Bessel sequences of unit norm vectors in a Hilbert space. Also, we prove that every Bessel sequence of unit vectors in a Hilbert space can be partitioned into finitely many uniformly separated sequences.

## 1. INTRODUCTION

There are many variations of the Feichtinger Conjecture, all equivalent with the following:

*Every Bessel sequence of unit vectors in a Hilbert space can be partitioned into finitely many Riesz sequences.*

For details on the Feichtinger Conjecture and the connection with other problems, see [1], [2], [3], [8], [9] and references of these papers.

We denote by  $\mathcal{H}$  a Hilbert space and  $\mathcal{F} = \{f_n\}_{n \in \mathbb{N}} \subset \mathcal{H}$ . We say that  $\mathcal{F}$  is a *Bessel sequence* if there exists  $B > 0$  so that

$$\sum_{n=0}^{\infty} |\langle x, f_n \rangle|^2 \leq B \|x\|^2, \quad \forall x \in \mathcal{H}.$$

$B$  is called Bessel constant for  $\mathcal{F}$ .

We say that  $\mathcal{F}$  is a *frame* for  $\mathcal{H}$  if it is a Bessel sequence and there exists  $A > 0$  so that

$$A \|x\|^2 \leq \sum_{n=0}^{\infty} |\langle x, f_n \rangle|^2, \quad \forall x \in \mathcal{H}.$$

For important applications of frames see the references of the paper [6].

We say that  $\mathcal{F}$  is a *Riesz sequence* (or *Riesz basic sequence*) if there are  $A, B > 0$  such that

$$A \sum |c_k|^2 \leq \left\| \sum c_k f_k \right\|^2 \leq B \sum |c_k|^2,$$

for any finite sequence  $(c_k)$ . Riesz sequences are particular cases of frames (see [4]).

---

2010 *Mathematics Subject Classification.* 46C05, 42C15.

*Key words and phrases.* Bessel sequence, Riesz sequence, Feichtinger Conjecture.

Let be  $I \subset \mathbb{N}$ . If  $\mathcal{F}$  is a Bessel sequences in  $\mathcal{H}$ , then  $\mathcal{F}_I = \{f_n\}_{n \in I}$  is clearly also a Bessel sequence in  $\mathcal{H}$ .

In [4], O. Christensen, using Schur's test, give conditions on a sequence  $\{f_n\}_{n=0}^\infty$  to be a Bessel sequence, that it only involves inner products between the elements  $\{f_n\}_{n=0}^\infty$ :

**Proposition 1.** [4] *Let  $\{f_n\}_{n=0}^\infty$  be a sequence in  $\mathcal{H}$  and assume that there exists a constant  $B > 0$  such that*

$$\sum_{k=0}^{\infty} |\langle f_j, f_k \rangle| \leq B, \quad \forall j \in \mathbb{N}.$$

*Then  $\{f_n\}_{n=0}^\infty$  is a Bessel sequence with bound  $B$ .*

We call this sequences *Bessel-Schur sequences*.

The intrinsically localized sequences, introduced by K. Gröchenig in [7], are particular cases of Bessel-Schur sequences. In the same paper, he proves that every localized frame is a finite union of Riesz sequences.

On the other hand, we recall the following definition:

**Definition 2.** [3] A sequence  $\{f_n\}_{n \in I}$  of unit vectors in  $\mathcal{H}$  is called *separated* if there exists a constant  $\gamma < 1$  such that

$$|\langle f_n, f_k \rangle| \leq \gamma$$

for any  $n, k \in \mathbb{N}$ ,  $n \neq k$ .

In [3], the authors, among others, give the following result:

**Theorem 3.** *Let  $\mathcal{H}$  be a Hilbert space and let  $\{f_n\}_{n \in I}$  be a Bessel sequence of unit vectors in  $\mathcal{H}$ . Then  $\{f_n\}_{n \in I}$  can be partitioned into finitely many separated Bessel sequences.*

In the following, we prove that the Bessel-Schur sequences satisfies the Feichtinger Conjecture. Also, we prove that every Bessel sequence of unit vectors in a Hilbert space can be partitioned into finitely many uniformly separated sequences.

## 2. THE RESULTS

First, we give a condition for a Bessel sequence of unit vectors to be a Riesz sequence.

**Theorem 4.** *Let  $\mathcal{F}_I = \{f_n\}_{n \in I}$  be a Bessel sequence of unit vectors. We suppose that*

$$\sigma := \sup_{j \in I} \sum_{\substack{i \in I \\ i \neq j}} |\langle f_i, f_j \rangle| < 1.$$

*Then  $\mathcal{F}_I$  is a Riesz sequence.*

*Proof.* If  $\mathcal{F}_I$  is a Bessel sequence in  $\mathcal{H}$ , then the following operators are linear and bounded:

$$T : l^2(I) \rightarrow \mathcal{H}, \quad T(c_i) = \sum_{i \in I} c_i f_i \quad (\text{synthesis operator})$$

$$\Theta : \mathcal{H} \rightarrow l^2(I), \quad \Theta x = \{\langle x, f_i \rangle\}_{i \in I} \quad (\text{analysis operator})$$

Moreover,  $\Theta$  is the adjoint of  $T$  (see [4]).

For  $c = (c_n)_{n \in I} \in l^2(I)$ , we have

$$\begin{aligned} (\Theta T)(c) &= \left\{ \left\langle \sum_{k \in I} c_k f_k, f_j \right\rangle \right\}_{j \in I} \\ &= \left\{ \sum_{k \in I} c_k \langle f_k, f_j \rangle \right\}_{j \in I} \end{aligned}$$

hence

$$(\Theta T)(c) - c = \left\{ \sum_{k \neq j} c_k \langle f_k, f_j \rangle \right\}_{j \in I}.$$

It follows, with Cauchy-Schwartz inequality,

$$\begin{aligned} \|(\Theta T)(c) - c\|_2^2 &= \sum_{j \in I} \left| \sum_{\substack{k \in I \\ k \neq j}} c_k \langle f_k, f_j \rangle \right|^2 \\ &\leq \sum_{j \in I} \left( \sum_{\substack{k \in I \\ k \neq j}} |c_k| |\langle f_k, f_j \rangle|^{1/2} \cdot |\langle f_k, f_j \rangle|^{1/2} \right)^2 \\ &\leq \sum_{j \in I} \left( \sum_{\substack{k \in I \\ k \neq j}} |c_k|^2 |\langle f_k, f_j \rangle| \right) \left( \sum_{\substack{k \in I \\ k \neq j}} |\langle f_k, f_j \rangle| \right) \\ &\leq \sigma \sum_{j \in I} \left( \sum_{\substack{k \in I \\ k \neq j}} |c_k|^2 |\langle f_k, f_j \rangle| \right) \end{aligned}$$

Changing the order of summation, we obtain

$$\begin{aligned} \|(\Theta T)(c) - c\|_2^2 &\leq \sigma \sum_{k \in I} |c_k|^2 \sum_{\substack{j \in I \\ j \neq k}} |\langle f_k, f_j \rangle| \\ &\leq \sigma^2 \|c\|_2^2, \end{aligned}$$

so  $\|\Theta T - I\| \leq \sigma < 1$ .

Therefore  $\Theta T$  is invertible, thus  $\Theta$  is surjective. It follows that  $\mathcal{F}_I$  is a Riesz-Fischer sequence. From Theorem 3 in [11], Ch. 4, Sec. 2, we have that there exists  $A > 0$  so that

$$A \sum |c_k|^2 \leq \left\| \sum c_k f_k \right\|^2$$

for every finite sequence  $(c_k)$ .

Since  $\mathcal{F}_I$  is a Bessel sequence, we have

$$\left\| \sum c_k f_k \right\|^2 \leq B \sum |c_k|^2$$

for  $(c_k)$  finite sequence (see [4]). So  $\mathcal{F}_I$  is a Riesz sequence.  $\square$

**Theorem 5.** *Every Bessel-Schur sequence of unit vectors is union of finite Riesz sequences.*

*Proof.* Let  $j \in \mathbb{N}$  fixed. We have:

$$\sum_{i=0}^{\infty} |\langle f_j, f_i \rangle| \leq B,$$

hence

$$(1) \quad \sum_{\substack{i=0 \\ i \neq j}} |\langle f_j, f_i \rangle| \leq B - 1, \quad \text{for any } j \in \mathbb{N}.$$

We denote

$$a_{ij} = \begin{cases} |\langle f_j, f_i \rangle| & , \quad j \neq i \\ 0 & , \quad j = i \end{cases}$$

We have  $a_{ij} = a_{ji} \geq 0$  and  $a_{ii} = 0$ .

The relation (1) is equivalent with

$$\sup_{j \in \mathbb{N}} \sum_{i \in \mathbb{N}} a_{ij} \leq B - 1.$$

By Mills' Lemma (see [5], Ch.X or [10]) there is a partition  $\mathbb{N} = I_1 \cup I_2$  such that

$$\sup_{j \in I_p} \sum_{i \in I_p} a_{ij} \leq \frac{B-1}{2}; \quad p = 1, 2.$$

By iteration, for any  $m \geq 1$ , there is a partition  $\mathbb{N} = I_1 \cup I_2 \cup \dots \cup I_{2^m}$  such that

$$\sup_{j \in I_p} \sum_{i \in I_p} a_{ij} \leq \frac{B-1}{2^m}, \quad \forall p = 1, 2, \dots, 2^m.$$

We take  $m$  so that  $\frac{B-1}{2^m} < 1$  and apply Theorem 4.  $\square$

### 3. AN EQUIVALENT FORM OF THE FEICHTINGER CONJECTURE

We consider the following class of sequences.

**Definition 6.** Let  $\mathcal{F}_I = \{f_n\}_{n \in I}$  be a sequence of unit vectors. We say that this sequence is *uniformly separated* if the following condition holds

$$\eta := \sup_{j \in I} \sum_{\substack{i \in I \\ i \neq j}} |\langle f_i, f_j \rangle|^2 < 1.$$

The following result is a refinement of a result from [3].

**Theorem 7.** *Every Bessel sequence of unit vectors is union of finite uniformly separated sequences.*

*Proof.* Let  $\mathcal{F}$  be a Bessel sequence of unit vectors:

$$\sum_{i=0}^{\infty} |\langle x, f_i \rangle|^2 \leq B \|x\|^2, \quad \forall x \in \mathcal{H}.$$

Let  $j \in \mathbb{N}$  fixed. We take  $x = f_j$ :

$$\sum_{i=0}^{\infty} |\langle f_j, f_i \rangle|^2 \leq B \|f_j\|^2 = B,$$

hence

$$(2) \quad \sum_{\substack{i=0 \\ i \neq j}}^{\infty} |\langle f_j, f_i \rangle|^2 \leq B - 1, \quad \text{for any } j \in \mathbb{N}.$$

It is clear that  $B \geq 1$ .

We denote

$$a_{ij} = \begin{cases} |\langle f_j, f_i \rangle|^2 & , \quad j \neq i \\ 0 & , \quad j = i \end{cases}$$

We have  $a_{ij} = a_{ji} \geq 0$  and  $a_{ii} = 0$ .

The relation (2) is equivalent with

$$\sup_{j \in \mathbb{N}} \sum_{i \in \mathbb{N}} a_{ij} \leq B - 1.$$

By Mills' Lemma (see [5], Ch.X or [10]) there is a partition  $\mathbb{N} = I_1 \cup I_2$  such that

$$\sup_{j \in I_p} \sum_{i \in I_p} a_{ij} \leq \frac{B-1}{2}; \quad p = 1, 2.$$

By iteration, for any  $m \geq 1$ , there is a partition  $\mathbb{N} = I_1 \cup I_2 \cup \dots \cup I_{2^m}$  such that

$$\sup_{j \in I_p} \sum_{i \in I_p} a_{ij} \leq \frac{B-1}{2^m}, \quad \forall p = 1, 2, \dots, 2^m.$$

We take  $m$  so that  $\frac{B-1}{2^m} < 1$  and apply Definition 6. □

From the above Theorem, we obtain the following equivalent form of the Feichtinger Conjecture:

*Every uniformly separated Bessel sequence of unit norm vectors can be partitioned into finitely many Riesz sequences.*

## REFERENCES

- [1] P.G. Casazza, O. Christensen, A.M. Lindner, R. Vershynin, Frames and the Feichtinger Conjecture, Proc. Amer. Math. Soc. 133 (4) (2005) 1025–1033.
- [2] P.G. Casazza, M. Fickus, J.C. Tremain, E. Weber, The Kadison-Singer problem in mathematics and engineering: A detailed account, Contemp. Math. 414 (2006) 299–355.
- [3] I. Chalendar, E. Fricain, D. Timotin, A short note on the Feichtinger Conjecture, arXiv:1106.3408v1, 2011.
- [4] O. Christensen, An Introduction to Frames and Riesz Bases, Birkhäuser, 2003.
- [5] J. Garnett, Bounded Analytic Functions, Academic Press, 1981.

- [6] P. Găvruta, On some identities and inequalities for frames in Hilbert spaces, J. Math. Anal. Appl. 321 (2006) 469–478.
- [7] K. Gröchenig, Localized frames are finite unions of Riesz sequences, Adv. Comput. Math. 18 (2003), 149–157.
- [8] S. Lata, V.I. Paulsen, Reproducing kernel Hilbert spaces, de Branges spaces and Feichtinger conjecture, Indiana Univ. Math. Journal 60(4)(2011)1303–1318.
- [9] S. Lata, The Feichtinger Conjecture and Reproducing Kernel Hilbert Spaces, PhD Thesis, University of Huston, 2010.
- [10] P. Thomas, Hardy spaces interpolation in the unit ball, Proc. Kon. Nederl. Akad. Wetensch. A-90. Indag. Math. 49 (1987) 325–351.
- [11] R. Young, An Introduction to Nonharmonic Fourier Series, Academic Press, New York, 1980.

”POLITEHNICA” UNIVERSITY OF TIMIȘOARA, DEPARTMENT OF MATHEMATICS, 300006, TIMIȘOARA, ROMANIA.

*E-mail address:* pgavruta@yahoo.com